



KENTUCKY

Children Aged <5 Years

2009

Pediatric Nutrition
Surveillance System

Graphics


Comparing Contributor and National Data

2008 (prior year) national PedNSS data are presented.

Contributors included 43 states, the District of Columbia, 6 Indian Tribal Organizations, and 2 U.S. territories.

Source of data

among children aged <5 years



* *Special Supplemental Nutrition Program for Women, Infants and Children.*


** *Early Periodic Screening, Diagnosis, and Treatment Program.*

+ *Title V Maternal and Child Health Program.*

++ *Includes Head Start.*


Racial and ethnic distribution

among children aged <5 years




Age distribution

among children aged <5 years



Income distribution*

among children aged <5 years




* Percent of poverty level.

** Adjunctive eligibility: eligible for Medicaid, Food Stamps, or other entitlement programs.

Program participation*

among children aged <5 years




* Represents program participation among children enrolled in the PH program contributing PedNSS data.

** Special Supplemental Nutrition Program for Women, Infants and Children.

*** Temporary Assistance for Needy Families.

Migrant status*


among children aged <5 years



* Family whose principal employment is agricultural on a seasonal basis; adapted from Federal Register 50744, 9/99.

Prevalence of low birthweight*

by race and ethnicity




* < 2500 grams, among infants born during the reporting period.

** Reduce low birthweight to 5% of live births.

Trends in prevalence of low birthweight*


by race and ethnicity



* < 2500 grams, among infants born during the reporting period.
Year 2010 target: reduce low birthweight to 5% of live births.

Prevalence of high birthweight*

by race and ethnicity




* > 4000 grams, among infants born during the reporting period.

2009 KY PedNSS Table 8C

Trends in prevalence of high birthweight*

by race and ethnicity




* > 4000 grams, among infants born during the reporting period.

2009 KY PedNSS Table 18C

Prevalence of short stature, underweight, and obesity*


among children aged <5 years



* Short stature: < 5th percentile length or height-for-age; underweight: < 5th percentile weight-for-length or BMI-for-age; obese: \geq 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.

Prevalence of short stature*

among children aged <5 years, by race and ethnicity




* < 5th percentile length or height-for-age, CDC Growth Charts, 2000.

** Reduce growth retardation (short stature) among low-income children under age five years to 5%. 2009 KY PedNSS Table 8C

Prevalence of short stature*

among children aged <5 years, by age




* < 5th percentile length or height-for-age, CDC Growth Charts, 2000.

** Reduce growth retardation (short stature) among low-income children under age five years to 5%.

2009 KY PedNSS Table 8C

Trends in prevalence of short stature*

among children aged <5 years, by race and ethnicity




* < 5th percentile length or height-for-age, CDC Growth Charts, 2000.

Year 2010 target: reduce growth retardation (short stature) among low-income children under age five years to 5%.

Prevalence of underweight*


among children aged <5 years, by race and ethnicity



* < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall below the 5th percentile.


Prevalence of underweight*

among children aged <5 years, by age



* < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall below the 5th percentile.


Trends in prevalence of underweight* among children aged <5 years, by race and ethnicity



* < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall below the 5th percentile.

Prevalence of obesity*


among children aged <5 years, by race and ethnicity



* ≥ 95 th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Prevalence of obesity*


among children aged <5 years, by age



* \geq 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Trends in prevalence of obesity*


among children aged <5 years, by race and ethnicity



* ≥ 95 th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Trends in prevalence of obesity*


among children aged <5 years, by age



* ≥ 95 th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Prevalence of obesity and overweight*


among children aged 2 to <5 years, by race and ethnicity



* Obese: \geq 95th percentile BMI-for-age; overweight: \geq 85th- <95th percentile BMI-for-age, CDC Growth Charts, 2000. 15% of children are expected to fall above the 85th percentile (5% above the 95th percentile and 10% between the 85th and 95th percentiles).

Prevalence of obesity and overweight*


among children aged 2 to <5 years, by age



* Obese: \geq 95th percentile BMI-for-age; overweight: \geq 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000. 15% of children are expected to fall above the 85th percentile (5% above the 95th percentile and 10% between the 85th and 95th percentiles).

Trends in prevalence of overweight*


among children aged 2 to <5 years, by race and ethnicity



* $\geq 85^{\text{th}}\text{--}<95^{\text{th}}$ percentile BMI-for-age, CDC Growth Charts, 2000.
10% of children are expected to fall between these percentiles.

Trends in prevalence of obesity*


among children aged 2 to <5 years, by race and ethnicity



* ≥ 95 th percentile BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Prevalence of anemia*


among children aged <5 years, by race and ethnicity



* Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

Prevalence of anemia*


among children aged <5 years, by age



* Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

Trends in prevalence of anemia*


among children aged <5 years, by race and ethnicity



* Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

Trends in prevalence of anemia*


among children aged <5 years, by age



* Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

Percentage of infants ever breastfed*

by race and ethnicity




* Among infants born during the reporting period.

** Increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

Trends in the percentage of infants ever breastfed*

by race and ethnicity




* Among infants born during the reporting period.

Year 2010 target: increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

Percentage of infants breastfed at least 6 months*

by race and ethnicity




* Among infants who turned six months of age during the reporting period.

** Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

Trends in the percentage of infants breastfed at least 6 months*

by race and ethnicity




* Among infants who turned six months of age during the reporting period.

Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

2009 KY PedNSS Table 19C

Percentage of infants breastfed at least 12 months*

by race and ethnicity




* Among infants who turned twelve months of age during the reporting period.

** Increase the proportion of mothers who breastfeed their babies at one year to 25%.

Trends in the percentage of infants breastfed at least 12 months*


by race and ethnicity



* Among infants who turned twelve months of age during the reporting period.

Year 2010 target: increase the proportion of mothers who breastfeed their babies at one year to 25%.

Trends in the percentage of infants ever breastfed, and breastfed at least 6 and 12 months




Year 2010 targets: increase the proportion of mothers who breastfeed their babies a) in the early postpartum period to 75%, b) at six months to 50%, and c) at one year to 25%.

Percentage of infants exclusively breastfed at least 3 and 6 months*

Insufficient Data


** Among infants who turned three and six months of age respectively during the reporting period.*

Percentage of children aged 2 to <5 years who view TV ≤ 2 hours/day by race and ethnicity



* Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.


Percentage of children aged 2 to <5 years who view TV ≤ 2 hours/day by age



* Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.

Percentage of children aged <5 years who live in households with smokers*

by race and ethnicity




* Defined as smokers who smoke inside the home.

** Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.

Percentage of children aged <5 years who live in households with smokers*

by age



* Defined as smokers who smoke inside the home.


** Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.

Graphics

Comparing Contributor and Local Data

Racial and ethnic distribution

among children aged <5 years, by region*




* Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 4E

Age distribution

among children aged <5 years, by region*




* Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 5E

Prevalence of low birthweight*

by region**




* < 2500 grams, among infants born during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

*** Reduce low birthweight to 5% of live births.

Prevalence of high birthweight*

by region**




* > 4000 grams, among infants born during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 6E

Prevalence of short stature*

among children aged <5 years, by region**




* < 5th percentile length or height-for-age, CDC Growth Charts, 2000.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

*** Reduce growth retardation (short stature) among low-income children under age five years to 5%. 2009 KY PedNSS Table 6E

Prevalence of underweight*

among children aged <5 years, by region**



* < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.


5% of children are expected to fall below the 5th percentile.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 6E

Prevalence of obesity*

among children aged <5 years, by region**




* \geq 95th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

Prevalence of obesity and overweight*

among children aged 2 to <5 years, by region**




* Obese: \geq 95th percentile BMI-for-age; overweight: \geq 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000. 5% of children are expected to fall above the 95th percentile; 15% of children are expected to fall above the 85th percentile.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 6E

Prevalence of anemia*

among children aged <5 years, by region**




* Hb or Hct < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

2009 KY PedNSS Table 6E

Percentage of infants ever breastfed*

by region**




* Among infants born during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

*** Increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

Percentage of infants breastfed at least 6 months*

by region**




* Among infants who turned six months of age during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

*** Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

Percentage of infants breastfed at least 12 months*

by region**



* Among infants who turned twelve months of age during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.

*** Increase the proportion of mothers who breastfeed their babies at one year to 25%.

Percentage of infants exclusively breastfed at least 3 months*


by region**

Insufficient Data

* Among infants who turned three months of age during the reporting period.

** Data for some regions not presented if <100 records are available for analysis after exclusions.


Percentage of children aged 2 to <5 years who view TV ≤ 2 hours/day by region*



* Data for some regions not presented if <100 records are available for analysis after exclusions.

** Increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.

Percentage of children aged <5 years who live in households with smokers* by region**



* Defined as smokers who smoke inside the home.

** Data for some regions not presented if <100 records are available for analysis after exclusions.


*** Reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.

Maps

State Maps of County Data

Prevalence of low birthweight*


by county



* < 2500 grams, among infants born during the reporting period.
Year 2010 target: reduce low birthweight to 5% of live births.

Prevalence of high birthweight*

by county




* > 4000 grams, among infants born during the reporting period.

2009 KY PedNSS Table 6B

Prevalence of short stature*

among children aged <5 years, by county




* < 5th percentile length or height-for-age, CDC Growth Charts, 2000.

Year 2010 target: reduce growth retardation (short stature) among low-income children under age five years to 5%.

Prevalence of underweight*


among children aged <5 years, by county



* < 5th percentile weight-for-length or BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall below the 5th percentile.

Prevalence of overweight*


among children aged 2 to <5 years, by county



* \geq 85th-<95th percentile BMI-for-age, CDC Growth Charts, 2000.
10% of children are expected to fall between the 85th and 95th percentiles.

Prevalence of obesity*


among children aged 2 to <5 years, by county



* \geq 95th percentile BMI-for-age, CDC Growth Charts, 2000.
5% of children are expected to fall above the 95th percentile.

Prevalence of anemia*


among children aged <5 years, by county



* *Hb or Hct* < 5th percentile, CDC MMWR vol. 47 (No. RR-3), 1998.

Percentage of infants ever breastfed*

by county



* Among infants born during the reporting period.

Year 2010 target: increase the proportion of mothers who breastfeed their babies in the early postpartum period to 75%.

2009 KY PedNSS Table 7B


Percentage of infants breastfed at least 6 months*, by county

Insufficient Data

** Among infants who turned six months of age during the reporting period.*

Year 2010 target: increase the proportion of mothers who breastfeed their babies at six months to 50%.

Percentage of infants breastfed at least 12 months*, by county



* Among infants who turned twelve months of age during the reporting period.

Year 2010 target: increase the proportion of mothers who breastfeed their babies at one year to 25%.


2009 KY PedNSS Table 7B

Percentage of infants exclusively breastfed at least 3 months*, by county

Insufficient Data


** Among infants who turned three months during the reporting period.*

Percentage of children aged 2 to <5 years who view TV \leq 2 hours/day, by county



Year 2010 target: increase the proportion of children and adolescents who view TV two or fewer hours per day to 75%.

Percentage of children aged <5 years who live in households with smokers*, by county



* Defined as smokers who smoke inside the home.

Year 2010 target: reduce the proportion of children who are regularly exposed to tobacco smoke at home to 10%.

2009 KY PedNSS Table 7B

